

Video Abstracts

Pseudodystonic Posture Secondary to Klippel–Feil Syndrome and Diastematomyelia

Martin Lopez-Vicchi¹, Gustavo Da Prat¹ & Emilia Mabel Gatto^{1,2*}

¹Sanatorio de la Trinidad Mitre, Buenos Aires, Argentina, ²Department of Movement Disorders, Instituto Neurociencias de Buenos Aires (INEBA), Buenos Aires, Argentina

Abstract

Background: Dystonic postures possess a great number of differential diagnoses.

Phenomenology Shown: We describe a pseudodystonic posture in a 61-year-old woman with skeletal and extra-skeletal abnormalities.

Educational Value: Klippel–Feil syndrome represents an unusual cause of pseudodystonic posture to be considered in the differential diagnosis of dystonia.

Keywords: Pseudodystonia, Klippel–Feil Syndrome, diastematomyelia

Citation: Lopez-Vicchi ML et al. Pseudodystonic posture secondary to Klippel–Feil syndrome and diastematomyelia. Tremor Other Hyperkinet Mov 2015; 5. doi: 10.7916/D8ZC820C

*To whom correspondence should be addressed. E-mail: emiliagatto@fibertel.com.ar; Ga_mem@yahoo.com

Editor: Elan D. Louis, Yale University, USA

Received: May 17, 2015 **Accepted:** June 4, 2015 **Published:** July 1, 2015

Copyright: © 2015 Lopez-Vicchi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution–Noncommercial–No Derivatives License, which permits the user to copy, distribute, and transmit the work provided that the original author(s) and source are credited; that no commercial use is made of the work; and that the work is not altered or transformed.

Funding: None.

Financial Disclosures: None.

Conflict of Interest: The authors report no conflict of interest.

A 61-year-old female born of full-term normal delivery after a non-consanguineous marriage, reported congenital short neck with abnormal posture, unilateral sensorineural hearing loss, and migraine. She sought consultation at our center for chronic headaches and reported that when she was a teenager she was misdiagnosed with cervical dystonia. Clinical examination showed head tilt, limitation of movement at the cervical level, and low posterior hair implantation (Video 1). Spinal magnetic resonance imaging revealed vertebral fusion between the medulla and C4 as well as partial splitting between C2 and C3.

Klippel–Feil syndrome (KFS) is a rare congenital disorder (1 in 50,000 people), characterized by a defect in the formation or segmentation of the cervical vertebrae with short neck, low posterior hairline, and limited neck movement.¹ Extra-skeletal manifestations and spinal abnormalities including posterior fossa cysts or diastematomyelia may occur in KFS.²

In this case, KFS represents an unusual mimicking dystonia or pseudodystonic syndrome where sustained muscle contractions, unusual twisting movements, and/or abnormal postures occur secondary to underlying abnormalities.



Video 1. Clinical Examination. Head tilt, posture-mimicking laterocollis (pseudodystonia) with low posterior hairline in Klippel–Feil syndrome.

Acknowledgments

We thank the patient for her participation.

References

1. Brokinkel B, Wiebe K, Hesselmann V, et al. Surgical treatment in a patient with Klippel-Feil syndrome and anterior cervical meningocele: A case report and review of literature. *Eur Spine J* 2013;22:S517–S520. <http://dx.doi.org/10.1007/s00586-013-2769-6>.
2. Dickerman RD, Colle KO, Mittler MA. Intramedullary inflammatory mass dorsal to the Klippel–Feil deformity: error in development or response to an abnormal motion segment? *Spinal Cord* 2004;42:720–722. <http://dx.doi.org/10.1038/sj.sc.3101645>.